

Barrow Neurological Institute at St. Joseph's Hospital and Medical Center

## Barrow - St. Joseph's Scholarly Commons

---

Translational Neuroscience

---

12-9-2010

### Nicotinic Acetylcholine Receptor Efficacy And Pharmacological Properties Of 3-(Substituted Phenyl)-2<sup>1</sup>-Substituted Tropanes

F. Ivy Carroll

Bruce E. Blough

S. Wayne Mascarella

Hernán A. Navarro

J. Brek Eaton

*See next page for additional authors*

Follow this and additional works at: <https://scholar.barrowneuro.org/neurobiology>

---

#### Recommended Citation

Carroll, F. Ivy; Blough, Bruce E.; Mascarella, S. Wayne; Navarro, Hernán A.; Eaton, J. Brek; Lukas, Ronald J.; and Damaj, M. Imad, "Nicotinic Acetylcholine Receptor Efficacy And Pharmacological Properties Of 3-(Substituted Phenyl)-2<sup>1</sup>-Substituted Tropanes" (2010). *Translational Neuroscience*. 256.  
<https://scholar.barrowneuro.org/neurobiology/256>

This Article is brought to you for free and open access by Barrow - St. Joseph's Scholarly Commons. It has been accepted for inclusion in Translational Neuroscience by an authorized administrator of Barrow - St. Joseph's Scholarly Commons. For more information, please contact [rogena.lake@commonspirit.org](mailto:rogena.lake@commonspirit.org), [lynda.orescanin@commonspirit.org](mailto:lynda.orescanin@commonspirit.org), [laura.repak@commonspirit.org](mailto:laura.repak@commonspirit.org), [nicole.galvan@commonspirit.org](mailto:nicole.galvan@commonspirit.org), [suefue.espe@commonspirit.org](mailto:suefue.espe@commonspirit.org).

---

**Authors**

F. Ivy Carroll, Bruce E. Blough, S. Wayne Mascarella, Hernn A. Navarro, J. Brek Eaton, Ronald J. Lukas, and M. Imad Damaj

## Nicotinic Acetylcholine Receptor Efficacy and Pharmacological Properties of 3-(Substituted phenyl)-2 $\beta$ -substituted Tropanes

F. Ivy Carroll,<sup>\*,†</sup> Bruce E. Blough,<sup>†</sup> S. Wayne Mascarella,<sup>†</sup> Hernán A. Navarro,<sup>†</sup> J. Brek Eaton,<sup>‡</sup> Ronald, J. Lukas,<sup>‡</sup> and M. Imad Damaj<sup>§</sup>


<sup>†</sup>Center for Organic and Medicinal Chemistry, Research Triangle Institute, Research Triangle Park, North Carolina 27709-2194, United States, <sup>‡</sup>Division of Neurobiology, Barrow Neurological Institute, 350 West Thomas Road, Phoenix, Arizona 85013, United States, and <sup>§</sup>Department of Pharmacology and Toxicology, Medical Campus, Virginia Commonwealth University, Richmond, Virginia 23219, United States

Received August 2, 2010

There is a need for different and better aids to tobacco product use cessation. Useful smoking cessation aids, bupropion (**2**) and varenicline (**3**), share some chemical features with 3-phenyltropanes (**4**), which have promise in cocaine dependence therapy. Here we report studies to generate and characterize pharmacodynamic features of 3-phenyltropane analogues. These studies extend our work on the multiple molecular target model for aids to smoking cessation. We identified several new 3-phenyltropane analogues that are superior to **2** in inhibition of dopamine, norepinephrine, and sometimes serotonin reuptake. All of these ligands also act as inhibitors of nicotinic acetylcholine receptor (nAChR) function with a selectivity profile that favors, like **2**, inhibition of  $\alpha 3\beta 4^*$ -nAChR. Many of these ligands also block acute effects of nicotine-induced antinociception, locomotor activity, and hypothermia. Importantly, all except one of the analogues tested have better potencies in inhibition of nicotine conditioned place preference than **2**. We have identified new compounds that have utility as research tools and possible promise for treatment of nicotine dependence.

### Introduction

Tobacco product use, principally through cigarette smoking, is the greatest preventable cause of premature mortality, contributing in the United States to over 435 000 deaths annually.<sup>1</sup> Tobacco use cessation can halt and reverse the biological damage caused by smoking.<sup>2</sup> It is now commonly accepted that smoking behavior is maintained by the reinforcing effects of nicotine (**1**) and aversive effects of nicotine withdrawal.<sup>3–6</sup> Both nonpharmacological and pharmacological interventions have demonstrated efficacy in smoking cessation.<sup>7</sup> At present, first-line pharmaceutical treatments include nicotine replacement therapy (NRT),<sup>8</sup> bupropion (**2**) and varenicline (**3**).<sup>9,10</sup> While these treatments are useful in helping 5–20% of smokers abstain over the long term, new pharmacotherapies are needed that are either more effective or can impact those individuals not helped by existing treatments.



\*To whom correspondence should be addressed. Telephone: 919-541-6679. Fax: 919-541-8868. E-mail: fic@rti.org.

<sup>a</sup> Abbreviations: NRT, nicotine replacement therapy; DA, dopamine; 5HT, serotonin; NE, norepinephrine; HEK, human embryonic kidney; DAT, dopamine transporter; SERT, serotonin transporter; NET, norepinephrine transporter; nAChR, nicotinic acetylcholine receptor; VTA, ventral tegmental area; MPE, maximum possible effect; CPP, conditioned place preference.

The effectiveness of varenicline in smoking cessation is thought to be due to its action as a partial agonist at  $\alpha 4\beta 2$ -containing nicotinic acetylcholine receptors (nAChR).<sup>9,10</sup> The mechanism for bupropion's effectiveness as a smoking-cessation aid appears to be more multifaceted.<sup>11</sup> Its behavioral and neurophysiological effects resemble those of psychomotor stimulants,<sup>12</sup> and similar to other psychomotor stimulants, **2** inhibits the reuptake of dopamine (DA).<sup>13,14</sup> It also inhibits the reuptake of norepinephrine (NE)<sup>13,14</sup> and is a noncompetitive inhibitor of  $\alpha 3\beta 4^*$ -nAChR (where the \* indicates that nAChR subunits are known or possible assembly partners in addition to those indicated) and  $\alpha 4\beta 2$ -nAChR.<sup>14</sup> In animal behavioral pharmacology studies, **2** induced locomotor activity,<sup>15,16</sup> generalized to cocaine and amphetamine in drug discrimination studies,<sup>17,18</sup> produced conditioned place preference (CPP),<sup>19</sup> and was self-administered by both rats<sup>20</sup> and nonhuman primates.<sup>21</sup>

Over the past several years, we have synthesized a large number of 3-phenyltropane analogues and evaluated them for binding at monoamine transporters.<sup>22,23</sup> Similar to **2**, some analogues were better at inhibiting the dopamine (DA) transporter (DAT) and the norepinephrine (NE) transporter (NET) than the serotonin (5HT) transporter (SERT). Others were selective for DAT relative to NET and SERT, whereas others showed similar inhibition at all three transporters.<sup>22,23</sup> In 1995, Lerner-Marmorosh et al. reported that a number of 3-phenyltropane analogues were effective in blocking nicotine-induced seizures in mice and that a good correlation was observed between pharmacological potencies and abilities to block [<sup>3</sup>H]mecamylamine binding to brain membranes.<sup>24</sup> On the basis of these results, Lerner-Marmorosh et al. concluded that

the 3-phenyltropane analogues are neuronal nicotinic antagonists acting on a similar site to that of mecamylamine, a noncompetitive nicotinic antagonist. In another study, we reported that several 3-phenyltropane analogues blocked nicotine-induced antinociception in the tail-flick test in mice with potencies greater than that of **2**.<sup>25</sup> These intriguing results suggested that an additional pharmacological study of the 3-phenyltropane class of monoamine uptake inhibitor might provide information about the mechanism of action of non-competitive nicotinic antagonists like **2** and could provide lead compounds for development as aids to smoking cessation or as drugs for treatment of neurological or psychiatric disorders involving nicotinic mechanisms.

In this study, we report the synthesis and biological evaluation of 3-phenyltropane analogues **4a–l**. All analogues show inhibitory potency at human DAT and NET and functional antagonism of human  $\alpha 3\beta 4^*$ -nAChR. Similar to **2**, the compounds antagonize the antinociceptive, hypolocomotor, and/or hypothermic effects induced by an acute injection of nicotine in mice and blocked nicotine CPP after repeated injection.

## Chemistry


The 3-phenyltropane analogues **4a–e**, **4g**, and **4l** were synthesized as previously reported.<sup>26–29</sup> Scheme 1 outlines the synthetic route used to prepare  $3\beta$ -(4-chlorophenyl)-2 $\beta$ -(4',5'-dimethylbenzimidazol-2'-yl)tropane (**4f**).  $3\beta$ -(4-chlorophenyl)tropane-2 $\beta$ -carboxylic acid (**5**)<sup>28</sup> is converted to the acid chloride and then treated with 4,5-dimethyl-1,2-phenylenediamine to give amide **6**. Treatment of **6** with phosphorus oxychloride gave the desired **4f**. The nortropane analogue **4h** was prepared from **4c** as shown in Scheme 2. Treatment of **4c** in 1,2-dichloroethane containing excess 1-chloroethyl chloroformate (ACE-Cl) followed by refluxing the intermediate urethane in methanol gave **4h**.

The synthesis of the  $3\beta$ -(4-chloro-3-methylphenyl)-2 $\beta$ -(3'-substituted isoxazol-5'-yl)tropanes (**4i** and **4j**) is also outlined in Scheme 2. A solution of **4c** was added to the dilithium salt of the appropriate ketone oxime in tetrahydrofuran (THF) at 0 °C under nitrogen, and the reaction mixture was allowed to warm to 25 °C. After a few hours, the reaction mixture was added to a THF solution containing sulfuric acid and refluxed for 1 h to give the desired products **4i** and **4j**.

The synthesis of the  $3\alpha,2\beta$ -tropane **4k** is outlined in Scheme 3. Addition of a solution of (1*R*,5*S*)-2-(3'-methyl-1',2',4'-oxadiazol-5'-yl)-8-methyl-8-azabicyclo[3.2.1]oct-2-ene (**7**)<sup>30</sup> in anhydrous THF at -78 °C to a solution of the 3-chloro-4-methylphenyllithium (prepared from the appropriate aryl bromide and butyllithium) followed by quenching with 1 N hydrochloric acid at -78 °C formed the  $3\alpha$ -(substituted phenyl)tropane-2 $\alpha$ -(3'-methyl-1',2',4'-oxadiazol-5'-yl)tropanes (**8**). In addition to the desired isomer, the  $2\alpha,3\beta$ -isomer was also formed, which was removed by flash chromatography or carried through and removed at the next reaction. Transformation of oxadiazole **8** to the desired methyl ester **4k** was accomplished by reduction with nickel boride (generated in situ by reaction of sodium borohydride and nickel tetraacetate) and hydrochloric acid in refluxing methanol. Under such conditions, a complete epimerization of C-2 to form the  $3\alpha,2\beta$ -stereoisomer was observed.


**In Vitro Assays.** The 3-phenyltropane analogues **4a–l** were evaluated for their ability to block reuptake of [<sup>3</sup>H]dopamine ([<sup>3</sup>H]DA), [<sup>3</sup>H]serotonin ([<sup>3</sup>H]SERT), and [<sup>3</sup>H]norepinephrine

## Scheme 1<sup>a</sup>




<sup>a</sup> Reagents: (a) (COCl)<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>; (b) 4,5-dimethyl-1,2-phenylenediamine, CH<sub>2</sub>Cl<sub>2</sub>; (c) POCl<sub>3</sub>.

## Scheme 2<sup>a</sup>



<sup>a</sup> Reagents: (a) ACE-Cl, DCE, reflux; (b) CH<sub>3</sub>OH, reflux; (c) CH<sub>3</sub>-(C=NOH)R, *n*-C<sub>4</sub>H<sub>9</sub>Li, THF, 0 °C; (d) H<sub>2</sub>SO<sub>4</sub>, THF.

## Scheme 3<sup>a</sup>



<sup>a</sup> Reagents: (a) *n*-C<sub>4</sub>H<sub>9</sub>Li, THF, -78 °C, 4 h; (b) TFA; (c) Ni<sub>2</sub>B, CH<sub>3</sub>OH, HCl.

([<sup>3</sup>H]NE) using human (h) DAT, hSERT, and hNET stably expressed in HEK293 cells using conditions similar to those previously reported.<sup>31,32</sup> The results are given in Table 1.


The 3-phenyltropane analogues **4a–l** were also evaluated for their ability to antagonize functional responses of  $\alpha 3\beta 4^*$ -,  $\alpha 4\beta 2$ -,  $\alpha 4\beta 4$ -, and  $\alpha 1^*$ -nAChR and mechanisms involved for functional inhibition using previously reported methods.<sup>33</sup> Results are given in Table 1 and in Figures 1 and 2.<sup>32</sup>

**In Vivo Assays. Acute Nicotine Testing.** The 3-phenyltropane analogues **4a–l** were also evaluated for their ability to antagonize behavioral responses to acute nicotine administration as previously described.<sup>32</sup> Results are given in Table 2.

**Nicotine Reward Using the CPP Test.** Selected 3-phenyltropane analogues were also evaluated for their ability to antagonize the development of nicotine-induced CPP in mice using an unbiased paradigm.<sup>34</sup> Results are given in Table 2.

## Results

**Effects on Monoamine Uptake.** Compound **2** is a relatively weak monoamine uptake inhibitor. In contrast, all of the 3-phenyltropane analogues studied here have higher inhibitory potencies for DA uptake inhibition.  $3\beta$ -(4-Methylphenyl)tropane-2 $\beta$ -carboxylic acid isopropyl ester (**4a**) and

**Table 1.** 3-Phenyltropane Analogue Inhibition of Monoamine Uptake and Nicotinic Acetylcholine Receptor (nAChR) Function

Compd	Stereo-chemistry	R <sub>1</sub>	R <sub>2</sub>	X	Y	Monoamine Uptake Inhibition IC <sub>50</sub> (nM) <sup>a</sup>			nAChR Inhibition IC <sub>50</sub> , (μM) <sup>b</sup>			
						[ <sup>3</sup> H]DA	[ <sup>3</sup> H]NE	[ <sup>3</sup> H]5HT	α3β4*-	α4β2-	α4β4-	α1*-
<b>2</b>						658 ± 180	1850 ± 300	IA	1.8 (1.15)	12 (1.15)	15 (1.07)	7.9 (1.12)
<b>4a</b>	2β,3β	CH <sub>3</sub>	CO <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	H	0.83 ± 0.20	52 ± 10	2830 ± 500	1.8 (1.07)	12 (1.15)	8.8 (1.13)	12 (1.07)
<b>4b</b>	2β,3β	CH <sub>3</sub>	CO <sub>2</sub> -	CH <sub>3</sub>	H	4.8 ± 0.5	122 ± 20	2010 ± 800	1.3 (1.06)	11 (1.12)	7.3 (1.15)	9.5 (1.05)
<b>4c</b>	2β,3β	CH <sub>3</sub>		CH <sub>3</sub>	H	0.75 ± 0.1	11 ± 2	2600 ± 1300	1.7 (1.05)	15 (1.10)	9.9 (1.11)	9.9 (1.05)
<b>4d</b>	2β,3β	CH <sub>3</sub>		CH <sub>3</sub>	H	11 ± 3	310 ± 130	IA	20 (1.12)	942 (1.74)	57 (4.47)	95 (1.06)
<b>4e</b>	2β,3β	CH <sub>3</sub>		Cl	H	3.9 ± 1	26 ± 7	2000 ± 300	1.8 (1.03)	15 (1.15)	>100	19 (1.03)
<b>4f</b>	2β,3β	CH <sub>3</sub>		Cl	H	7.3 ± 2	19 ± 4	IA	0.57 (1.03)	1.8 (1.15)	4.3 (1.03)	4.7 (1.06)
<b>4g</b>	2β,3β	CH <sub>3</sub>	CO <sub>2</sub> CH <sub>3</sub>	Cl	CH <sub>3</sub>	2 ± 1	1.1 ± 0.3	1.0 ± 0.3	3.6 (1.10)	9.8 (1.18)	24 (1.16)	26 (1.07)
<b>4h</b>	2β,3β	H	CO <sub>2</sub> CH <sub>3</sub>	Cl	CH <sub>3</sub>	2.3 ± 0.4	0.9 ± 0.3	6.2 ± 2	3.1 (1.12)	16 (1.06)	12 (1.04)	11 (1.07)
<b>4i</b>	2β,3β	CH <sub>3</sub>		Cl	CH <sub>3</sub>	1.5 ± 0.3	6.3 ± 0.9	115 ± 20	0.73 (1.12)	7.7 (1.26)	5.8 (1.08)	13 (1.18)
<b>4j</b>	2β,3β	CH <sub>3</sub>		Cl	CH <sub>3</sub>	15 ± 5	14 ± 3	700 ± 200	1.4 (1.19)	14 (1.06)	4.6 (1.04)	3.0 (1.09)
<b>4k</b>	2β,3α	CH <sub>3</sub>	CO <sub>2</sub> CH <sub>3</sub>	Cl	CH <sub>3</sub>	3 ± 1	6.5 ± 2	105 ± 1	3.2 (1.08)	29 (1.04)	15 (1.04)	13 (1.05)
<b>4l</b>	2β,3α	H	CO <sub>2</sub> CH <sub>3</sub>	CH <sub>3</sub>	F	6.4 ± 2	0.5 ± 0.1	110 ± 7	2.6 (1.07)	11 (1.12)	25 (1.36)	7.3 (1.04)


<sup>a</sup> Values for mean ± standard error of three independent experiments, each conducted with triplicate determination. <sup>b</sup> Mean micromolar IC<sub>50</sub> values (to two significant digits) for bupropion and the indicated 3-phenyltropane analogues from three independent experiments for inhibition of functional responses to an EC<sub>80</sub>–EC<sub>90</sub> concentration of carbamylcholine mediated by nAChR subtypes composed of the indicated subunits (where \* indicates that additional subunits are or may be additional assembly partners with the subunits specified; see Experimental Section). Numbers in parentheses indicate SEM as a multiplication/division factor of the mean micromolar IC<sub>50</sub> values shown [i.e., the value 1.8 (1.15) reflects a mean IC<sub>50</sub> value of 1.8 μM with an SEM range of 1.8 × 1.15 μM to 1.8/1.15 μM or 1.6–2.1 μM]. IA: IC<sub>50</sub> > 100 μM.

3β-(4-methylphenyl)-2β-(3'-ethylisoxazol-5'-yl)tropane (**4c**), with IC<sub>50</sub> values of 0.83 and 0.75 nM, respectively, were the most potent analogues as inhibitors of DA uptake. However, 3β-(4-chloro-3-methylphenyl)-2β-(3'-methylisoxazol-5'-yl)tropane (**4i**), 3β-(4-chlorophenyl-3-methyl)tropane-2β-carboxylic acid methyl ester (**4g**), and its nortropane analogue (**4h**), with IC<sub>50</sub> values of 1.5, 2.0, and 2.3 nM, are almost as potent at inhibition of DA uptake as **4a** and **4c**. In addition, all of the 3-phenyltropane analogues were potent at NE uptake inhibitors. The most potent NE uptake inhibitors were the nortropane analogues 3α-(4-fluoro-3-methylphenyl)nortropane-2β-carboxylic acid methyl ester (**4l**) and 3β-(4-chloro-3-methylphenyl)-nortropane-2β-carboxylic acid methyl ester (**4h**), with IC<sub>50</sub> values of 0.5 and 0.9 nM, respectively. 3α-(4-Chloro-3-methylphenyl)tropane-2β-carboxylic acid methyl ester (**4g**), 3β-(4-chloro-3-methylphenyl)-2β-(3'-methylisoxazol-5'-yl)tropane (**4i**),


and 3α-(4-chloro-4-methylphenyl)tropane 2β-carboxylic acid methyl ester (**4k**), with IC<sub>50</sub> values of 1.1, 6.3, and 6.5, respectively, also are very potent NE uptake inhibitors.

Most of the 3-phenyltropane analogues are inactive or have IC<sub>50</sub> values greater than 100 nM for 5HT uptake inhibition. However, the 4-chloro-3-methylcarboxylic acid methyl ester **4g** and its nortropane analogue **4h** have IC<sub>50</sub> values of 1 and 6.2 nM for 5HT uptake inhibition. Since these two analogues also have high potency uptake inhibition at all three monoamine transporters.

Analogue **4g** has a slight (~2-fold) preference for inhibition of 5HT and NE over DA uptake inhibition. Compound **4j** has comparable inhibitory potencies for DA and NE uptake. Ligand **4h** has slight (~2-fold) preference for NE over DA uptake inhibition. Compound **4l** has 14-fold preference for



**Figure 1.** Specific  $^{86}\text{Rb}^+$  efflux (ordinate, percentage of control) was determined for functional, human muscle-type  $\alpha 1\beta 1\gamma\delta$ -nAChR ( $\Delta$ ), ganglionic  $\alpha 3\beta 4^*$ -nAChR ( $\blacktriangledown$ ),  $\alpha 4\beta 2$ -nAChR ( $\circ$ ), or  $\alpha 4\beta 4$ -nAChR ( $\blacktriangle$ ) naturally or heterologously expressed in human cell lines in the presence of a receptor subtype-specific,  $\text{EC}_{80}$ – $\text{EC}_{90}$  concentration of the full agonist, carbamylcholine, either alone or in the presence of the indicated concentrations (abscissa, log molar) of 3-phenyltropane analogue **4b**, **4c**, **4e**, or **4f** as indicated. Mean micromolar  $\text{IC}_{50}$  values and SEM as a multiplication/division factor of the mean micromolar  $\text{IC}_{50}$  value are provided in Table 1.



**Figure 2.** Specific  $^{86}\text{Rb}^+$  efflux (ordinate, percentage of control) was determined for functional, human ganglionic  $\alpha 3\beta 4^*$ -nAChR naturally expressed by SH-SY5Y human neuroblastoma cells in the presence of the full agonist, carbamylcholine, at the indicated concentrations (abscissa, log molar) either alone ( $\blacksquare$ ) or in the presence of 10  $\mu\text{M}$  3-phenyltropane analogues **4b** ( $\Delta$ ), **4c** ( $\square$ ), **4e** ( $\blacktriangledown$ ), or **4f** ( $\bullet$ ) as indicated. The reduction of apparent agonist efficacy without an effect on potency is consistent with a noncompetitive mechanism for nAChR block.

inhibition of NE over DA uptake. Otherwise, all the other compounds have preference for inhibition of DA uptake over other monoamine transporters, as low as 2-fold for **4k** and **4f** to  $> 60$ -fold for **4a**.

**Effects on nAChR Function.** The effects of 3-phenyltropane analogues **4a**–**l** on function of diverse human nAChR subtypes naturally or heterologously expressed by human cell lines were assessed using  $^{86}\text{Rb}^+$  efflux assays that are specific only for nAChR function in the cells used. None of the analogues has activity as agonists at  $\alpha 1^*$ -,  $\alpha 3\beta 4^*$ -,  $\alpha 4\beta 2$ -, or  $\alpha 4\beta 4$ -nAChR because  $^{86}\text{Rb}^+$  efflux in the presence of these ligands alone at concentrations from  $\sim 5$  nM to 100  $\mu\text{M}$  (data not shown here) was indistinguishable from responses in cells exposed only to efflux buffer.

$^{86}\text{Rb}^+$  efflux assays also were used to assess whether ligands had activity as antagonists at human nAChR. Representative

concentration-response curves for selected ligands (Figure 1) illustrate nAChR in vitro inhibitory profiles (see also Table 1).

Compound **2** has  $\text{IC}_{50}$  values of 1.8, 12, 15, and 7.9  $\mu\text{M}$  for functional antagonism of  $\alpha 3\beta 4^*$ -,  $\alpha 4\beta 2$ -,  $\alpha 4\beta 4$ -, and  $\alpha 1\beta 1^*$ -nAChR, respectively. Analogues **4a**, **4b**, **4c**, **4e**, **4f**, **4i**, and **4j** all have  $\text{IC}_{50}$  values equal to or lower than that for **2** at  $\alpha 3\beta 4^*$ -nAChR. The most potent inhibitors of  $\alpha 3\beta 4^*$ -nAChR function are **4f** and **4i** ( $\text{IC}_{50} = 0.57$  and 0.73  $\mu\text{M}$ , respectively). Similar to **2**, all the 3-phenyltropane analogues show preference for functional inhibition of  $\alpha 3\beta 4^*$ -nAChR over the other nAChR subtypes tested, with greatest overall preference for **4e** (8-,  $> 56$ -, and 11-fold over  $\alpha 4\beta 2$ -,  $\alpha 4\beta 4$ -, or  $\alpha 1^*$ -nAChR) and **4i** (11-, 8-, and 18-fold over  $\alpha 4\beta 2$ -,  $\alpha 4\beta 4$ -, or  $\alpha 1^*$ -nAChRs). Only **4f** had significantly better potency as a functional antagonist of  $\alpha 4\beta 2$ -nAChR than **2**. Analogue **4j** had the lowest overall preference for  $\alpha 3\beta 4^*$ -nAChR ( $\sim 2$ -fold), in part because it (and **4f**) had the highest inhibitory potencies at  $\alpha 4\beta 4$ - and  $\alpha 1^*$ -nAChR. Compounds **4f** and **4g** had the lowest preference ( $\sim 3$ -fold), and compounds **4i** and **4j** had the highest preference (10- to 13-fold) for  $\alpha 3\beta 4^*$ - over  $\alpha 4\beta 2$ -nAChR.

As was the case for **2**, all of the 3-phenyltropane analogues tested inhibited nAChR function via an apparently noncompetitive mechanism, lowering apparent agonist efficacy without altering agonist  $\text{EC}_{50}$  values [representative data for compounds acting at  $\alpha 3\beta 4^*$ -nAChR are shown (Figure 2)].

Comparing inhibitory potencies across classes of targets, **4g**, **4h**, and **4i** have 3500- to 5700-fold selectivity for inhibition of NE uptake over  $\alpha 3\beta 4^*$ -nAChR, and selectivity for inhibition of DA uptake over  $\alpha 3\beta 4^*$ -nAChR ranges from 78- to 100-fold for **4f** and **4j** to  $> 2000$ -fold for **4a** and **4c**.

**In Vivo Effects.** Compound **2** blocks nicotine-induced antinociception in the tail-flick and hot-plate tests with  $\text{AD}_{50}$  values of 1.2 and 15 mg/kg, respectively. Ten of the 3-phenyltropane analogues were more potent in blocking nicotine's effects in the tail-flick assay than **2**, having  $\text{AD}_{50}$  values between 0.002 and 0.28 mg/kg. The most potent analogue in the tail-flick test was 3 $\alpha$ -(4-fluoro-3-methylphenyl)-2 $\beta$ -carboxylic acid methyl ester (**4l**),

**Table 2.** Comparison of Behavioral Potency for 3-Phenyltropane Analogues in Blocking Acute Effects of Nicotine (Antinociception, Hypomotility, and Hypothermia) and Development of Nicotine CPP<sup>a</sup>

compd	AD <sub>50</sub> (mg/kg)				
	tail-flick	hot-plate	locomotion	hypothermia	CPP
<b>2</b>	1.2 (1–1.8)	15 (6–19)	4.9 (0.9–46)	9.2 (4–23)	0.35
<b>4a</b>	0.015 (0.001–0.16)	IA	1.5 (0.3–7)	1.6 (0.3–9)	NT
<b>4b</b>	0.28 (0.09–0.64)	9.8 (0.7–34)	1 (0.1–2.1)	7 (2.5–19)	NT
<b>4c</b>	0.008 (0.001–0.04)	1.6 (0.7–3.4)	0.2 (0.1–2.1)	6.9 (2.5–19)	0.045
<b>4d</b>	IA	IA	8.7 (6.5–11.2)	IA	0.013
<b>4e</b>	0.018 (0.002–0.16)	IA	IA	IA	NT
<b>4f</b>	0.11 (0.04–0.3)	19 (5.2–70)	IA	IA	0.45
<b>4g</b>	0.21 (0.06–2.7)	0.35 (0.1–1.2)	0.11 (0.03–0.3)	0.14 (0.1–0.18)	0.25
<b>4h</b>	0.003 (0.001–0.01)	0.78 (0.05–12)	0.41 (0.1–1.7)	0.23 (0.1–4)	0.25
<b>4i</b>	0.009 (0.002–0.03)	3.3 (1.4–7.8)	0.37 (0.06–1.5)	0.61 (0.24–1.5)	0.09
<b>4j</b>	2.5 (1.3–5.8)	IA	IA	IA	0.07
<b>4k</b>	0.026 (0.004–0.1)	IA	5.3 (0.5–48)	7 (3.3–18)	0.15
<b>4l</b>	0.002 (0.001–0.06)	2.6 (0.6–16)	1.65 (0.7–3.9)	7.4 (0.8–14.2)	0.03

<sup>a</sup> Results for nicotine acute effects were expressed as AD<sub>50</sub> (mg/kg) ± confidence limits (CL). Dose-response curves were determined using a minimum of four different doses of test compound, and at least eight mice were used per dose group. AD<sub>50</sub> (mg/kg) values were estimated in the CPP test using a minimum of three different doses of test compound, and at least eight mice were used per dose group. IA: AD<sub>50</sub> > 20 mg/kg. NT: not tested.

AD<sub>50</sub> = 0.002 mg/kg. Six analogues, **4b**, **4c**, **4g**, **4h**, **4i**, and **4l**, with AD<sub>50</sub> values of 0.35–9.8 mg/kg were more potent than **2** in blocking nicotine's acute effects in the hot-plate test, with **4g** being most potent (AD<sub>50</sub> = 0.35 mg/kg, ~42-fold more potent than **2**).

Compound **2** blocks nicotine-induced increases in locomotor activity with an AD<sub>50</sub> value of 4.9 mg/kg. Compounds **4a**, **4b**, **4c**, **4g**, **4h**, **4i**, and **4l** with AD<sub>50</sub> values of 0.11–1.65 mg/kg are more potent than **2**. The most potent analogue was **4g**, which has an AD<sub>50</sub> value of 0.11 mg/kg.

Nicotine-induced hypothermia is blocked by **2** with an AD<sub>50</sub> of 9.2 mg/kg. Compounds **4a**, **4b**, **4c**, **4g**, **4h**, **4i**, **4k**, and **4l** with AD<sub>50</sub> values of 0.14–7.4 mg/kg are all more potent than **2**. Compound **4g** and its nortropane analogue **4h** with IC<sub>50</sub> values of 0.14 and 0.23 mg/kg, respectively, were the two most potent analogues in the hypothermia test.

Pretreatment with **2** at different doses decreased the development of nicotine-induced CPP in mice conditioned with 0.5 mg/kg nicotine, with an estimated AD<sub>50</sub> value of 0.35 mg/kg. Nine of the 3-phenyltropane analogues were tested for their ability to block nicotine's preference. Six of the compounds, **4c**, **4d**, **4i**, **4j**, **4k**, and **4l**, with AD<sub>50</sub> values of 0.013–0.15 mg/kg are more potent than **2**.

## Discussion

The results from this study show that the 3-phenyltropane analogues have monoamine uptake inhibition and nAChR antagonism profiles similar to those of **2**. Some of the analogues also have slightly higher potency as antagonists of  $\alpha 3\beta 4^*$ -nAChR, and all analogues retain preference across nAChR subtypes for blockade of  $\alpha 3\beta 4^*$ -nAChR. Moreover, some of these compounds show better potency than **2** as inhibitors of acute effects of nicotine and nicotine-induced CPP, which measures the acute rewarding effect of the drug.

Regardless of the type of substituent at the 2 $\beta$ - and 3 $\beta$ -position, all the 3-phenyltropane analogues (see structures **4a–l**) had high potency in DA uptake inhibition. In addition all the analogues except the 3 $\beta$ -cyclobutyl ester **4b** and the 3 $\beta$ -4-chlorophenylisoxazole **4d** also had high potency in NE uptake inhibition. From a structural perspective, extensive modifications of the 2 $\beta$ -group (see structures **4a–l**) lowered analogue potency as 5HT uptake inhibitors more than they altered inhibitory potency as DA and NE uptake inhibitors.

Only analogues with smaller 2 $\beta$ -substituents (**4g** and **4h**) retained high efficacy for 5HT uptake inhibition. On the basis of the limited comparisons, stereochemistry seems to influence activity for 5HT and NE uptake inhibition more than for DA uptake inhibition and  $\alpha 3\beta 4^*$ -nAChR antagonist potency (compare **4g** and **4k**). With the exception of the 3 $\beta$ -4-chlorophenylisoxazole analogue **4d**, all the analogues had similar relative potencies at  $\alpha 3\beta 4^*$ - and  $\alpha 4\beta 2$ -nAChRs.

It is interesting to note that analogue **4c** has 150 times more potency than **2** as an inhibitor of nicotine's analgesic action in the tail-flick assay, 9-fold better activity in the hot-plate assay, 24-fold better activity in hypolocomotion studies, and a marginally more potent effect on body temperature, whereas it has ~8-fold better activity in the CPP assay. Analogue **4c** does not differ much from **2** in its antagonistic potency at any of the nAChR subtypes studied, and it has 480-fold better activity as a DA uptake inhibitor, ~170-fold better activity as NE uptake inhibitor, and no less than 5-fold better activity as a 5HT uptake inhibitor than **2**. Since analogue **4g** (AD<sub>50</sub> = 0.25 mg/kg) has activity comparable to that of **2** (AD<sub>50</sub> = 0.35 mg/kg) in the CPP assay, is 6-fold more potent than **2** in the tail-flick assay, 40-, 44-, and 65-fold more potent than **2** in the hot-plate locomotor and hypothermia tests, and has about the same nAChR inhibition profile as **2** while having ~300 ~2000 > 10000 times more potency at DA/NE/5HT uptake inhibition, its CPP effects appear to be more related to antagonism of nAChR. A similar analysis of **4i** and **4j** shows that the 4- to 5-fold better CPP activity relative to **2** correlates better with their nAChR inhibition profile than with their monoamine transporter and acute nicotinic effects. Compound **4d** with AD<sub>50</sub> = 0.013 mg/kg was the most potent analogue in the CPP test and thus was 27 times more potent than **2**. Somewhat surprisingly, unlike **2**, compound **4d** was inactive in the acute mouse tail-flick, hot-plate, and hypothermia tests and less potent than **2** in the hypolocomotor test. However, it retained high antagonist potency against  $\alpha 3\beta 4^*$ -nAChR and in DA uptake inhibition. In previous studies, we found that even though **4d** was a potent and selective inhibitor of the DAT relative to the NET and SERT, it did not show locomotor activity in rats even though *in vivo* binding studies showed that appreciable brain levels were achieved.<sup>8,26</sup> We also found that **4d** was a positive allosteric modulator of the CB-1 cannabinoid receptor and suggested that its interaction with this receptor might in part be the reason for its unusual

behavioral pharmacological properties.<sup>8</sup> However, the reason for this lack of locomotor activity is still not fully understood. One possibility is that **4d** is interacting with some as yet unidentified target, possibly a nAChR subtype other than those studied, that is responsible for its lack of locomotor activity and potent activity in the CPP test.

Although many analogues have much better potency in behavioral assays than **2**, it is challenging to determine how those effects relate to ligand actions at molecular targets, in part because improvements over **2** for a given ligand can vary quite widely depending on the behavioral assay. Correlation plots comparing the *in vitro* and *in vivo* data did not provide any useful insights. Some of these challenges may be due to metabolism of analogues to more or less active forms. In addition, **2** in mice is converted in humans to hydroxy metabolites, one of which expresses much of the drug's behavioral activity.<sup>35</sup>

In summary, ligands have been developed that can serve as useful research tools, having different inhibitory potency profiles across nAChR and monoamine transporter targets. For example, *3β*-(4-methylphenyl)-2β-(3'-ethylisoxazol-5'-yl)-tropane (**4c**) has IC<sub>50</sub> values of 0.75, 11, and 2600 nM for inhibition of DA, NE, and 5HT uptake and an IC<sub>50</sub> of 1.7 μM for antagonism of α3β4\*-nAChR. Like **2**, **4c** is active in all four acute tests of nicotine action in mice. Importantly, it is 150, 9, and 5 times more potent in the tail-flick, hot-plate, and locomotor tests, respectively, than **2**. It also is 8 times more potent than **2** in blocking nicotine-CPP. Three other compounds (**4g**, **4h**, **4i**) have higher potencies than **4c** as inhibitors in all four tests of acute nicotine effects, and these compounds as well as **4d**, **4j**, **4k**, and **4l** have potency as antagonists of nicotine-CPP significantly higher than that of **2**. Compounds **4d** and **4j** are inactive or are less active than **2** in the acute test of nicotinic action. However, they retain high antagonist potency against α3β4\*-nAChR and DA uptake inhibition. The current studies support the idea that nAChR antagonism, particularly inhibition of α3β4\*-nAChR function, and inhibitory actions at monoamine transporters are pharmacodynamic features of 3-phenyltropanes. Perhaps most importantly, and certainly warranting further investigation, is the possible utility of these ligands as aids to smoking cessation, especially given their ability to inhibit acute effects of nicotine and a test of nicotine preference with better potencies than **2**, which is a useful pharmacotherapy for nicotine dependence.

## Experimental Section

Nuclear magnetic resonance (<sup>1</sup>H NMR and <sup>13</sup>C NMR) spectra were recorded on a 300 MHz (Bruker AVANCE 300) spectrometer. Chemical shift data for the proton resonances were reported in parts per million (δ) relative to internal (CH<sub>3</sub>)<sub>4</sub>Si (δ 0.0). Elemental analyses were performed by Atlantic Microlab, Norcross, GA. Purity of compounds (> 95%) was established by elemental analysis. Analytical thin-layer chromatography (TLC) was carried out on plates precoated with silica gel GHLF (250 μM thickness). TLC visualization was accomplished with a UV lamp or in an iodine chamber. All moisture-sensitive reactions were performed under a positive pressure of nitrogen maintained by a direct line from a nitrogen source. Anhydrous solvents were purchased from Aldrich Chemical Co. or VWR. CMA80 is a mixture of 80% chloroform, 18% methanol, and 2% concentrated ammonium hydroxide.

**3β**-(4-Chlorophenyl)-2β-(4',5'-dimethylbenzimidazol-2'-yl)tropane (**4f**) Dihydrochloride. Compound **6** (3.8 g, 0.0095 mol) and POCl<sub>3</sub> (50 mL) were mixed and stirred at reflux for 90 min. The reaction mixture was cooled and added to petroleum

ether (2 L). Most of the solvent was decanted, and concentrated NH<sub>4</sub>OH-water (1:1) was added until basic to litmus paper and then extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to yield 3.0 g of a tan amorphous solid. This solid was chromatographed on silica gel, eluting with EtOAc and then EtOAc-CMA80 (1:1) to afford 1.7 g (47%) of **4f**. The free base was dissolved in ether and treated with 1 M ethereal HCl to give 1.65 g of the dihydrochloride salt of **4f** as a white solid: mp 224–227 °C; [α]<sub>D</sub> -166.2 °C (*c* 0.69, MeOH). <sup>1</sup>H NMR (CDCl<sub>3</sub>, free base) δ 1.73 (bd, 1H), 1.88 (q, *J* = 9.0 Hz, 2H), 2.22–2.35 (m, 4H), 2.31(s, 3H), 2.33 (s, 3H), 2.35 (s, 3H), 3.20–3.48 (m, 4H), 6.67 (d, *J* = 9.0 Hz, 2H), 7.00 (d, *J* = 9.0 Hz, 2H), 7.26 (s, 2H). Anal. (C<sub>23</sub>H<sub>28</sub>Cl<sub>3</sub>N<sub>3</sub>·1.25H<sub>2</sub>O) C, H, N.

(-)-*N*-Nor-3β-(3-methyl-4-chlorophenyl)tropane-2β-carboxylic Acid Methyl Ester (**4h**) Tartrate. Compound **4c** (0.85 g, 2.76 mmol) was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (30 mL), and 1-chloroethyl chloroformate (ACE-Cl, 4.2 mL, 39 mmol) was added. The mixture was refluxed for 8 h. The reaction mixture was concentrated, the residue dissolved in MeOH (30 mL), and the solution refluxed overnight. The MeOH solution was concentrated and the residue partitioned between CH<sub>2</sub>Cl<sub>2</sub> and 25 mL of NH<sub>4</sub>OH-H<sub>2</sub>O (1:1). The layers were separated and the aqueous layer extracted twice with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to give 0.85 g of yellow solid, which was chromatographed on 50 g of silica gel using 25% CMA80 in CH<sub>2</sub>Cl<sub>2</sub> to obtain 0.54 g (67%) of **4h**. This material was converted to the tartrate salt by dissolving 520 mg (1.77 mmol) of **4h** in MeOH and adding a MeOH solution of *D*-tartaric acid 0.286 mg (1.77 mmol). The salt was crystallized from MeOH-ether to give 0.70 g of **4h**·tartrate: mp 156–158 °C; [α]<sub>D</sub><sup>25</sup> -100.0 °C (*c* 1.00, MeOH). Anal. Calcd for C<sub>20</sub>H<sub>25</sub>ClNO<sub>8</sub>: C, 54.12; H, 5.90; N, 3.16; Cl, 7.99. Found: C, 53.88; H, 6.01; N, 3.13; Cl, 7.88. <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.30 (d, 1H), 7.18 (s, 1H), 7.03 (d, 1H), 4.40 (s, 2H), 4.22 (m, 2H), 3.50 (m, 1H), 3.38 (s, 3H), 3.05 (d, 1H), 2.61 (t, 1H), 2.35 (s, 3H), 2.12–2.28 (m, 4H), 1.91 (d, 1H). Anal. (C<sub>20</sub>H<sub>26</sub>ClNO<sub>8</sub>) C, H, N.

(-)-3β-(3-Methyl-4-chlorophenyl)tropane-2β-(3'-methylisoxazol-5'-yl) (**4i**) Hydrochloride. Acetone oxime (262 mg, 359 mmol) was dissolved under N<sub>2</sub> in 7.5 mL of THF, and the solution was cooled to 0 °C. Butyllithium, 25 M in hexanes (2.9 mL, 7.18 mmol), was slowly added, and the solution was stirred in an ice bath for 2–3 h. Compound **4c** (85 mg, 2.76 mmol) was dissolved in 4 mL of THF and slowly added to the cold reaction mixture. The mixture was allowed to warm to room temperature and remained there for 16 h. A solution of 1.65 g of 36 N H<sub>2</sub>SO<sub>4</sub>, 2.1 mL of H<sub>2</sub>O, and 7.65 mL of THF was slowly added, and the mixture was refluxed for 6 h before being stirred at room temperature for 16 h. The reaction mixture was concentrated and basified with NH<sub>4</sub>OH-H<sub>2</sub>O (1:1) to pH 10–11 and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 10 mL). The combined organic extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated to give 0.87 g of product, which was purified by flash chromatography on silica gel using 10% Et<sub>2</sub>O-Et<sub>3</sub>N (9:1) in 90% hexanes to give 0.26 g of **4c** and 0.27 g (44% adjusted) of the desired **4i**. The product was converted to the HCl salt by adding 1 M HCl in Et<sub>2</sub>O to an Et<sub>2</sub>O solution of **4i**. Recrystallization from MeOH-Et<sub>2</sub>O gave **4i**·HCl: mp 150 °C (dec); [α]<sub>D</sub><sup>25</sup> -112.0 °C (*c* 1.00, MeOH). <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 7.20 (d, 1H), 7.08 (s, 1H), 6.94 (d, 1H), 5.70, (s, 1H), 4.22 (m, 1H), 4.12 (m, 1H), 3.87 (m, 1H), 3.68 (m, 1H), 2.89 (s, 3H), 2.16–2.66 (m, 16H). Anal. (C<sub>19</sub>H<sub>25</sub>Cl<sub>2</sub>N<sub>2</sub>O) C, H, N.

(-)-3β-(3-Methyl-4-chlorophenyl)tropane-2β-(3'-phenylisoxazol-5'-yl) (**4j**) Hydrochloride. Acetophenone oxime (617 mg, 4.56 mmol) was dissolved under N<sub>2</sub> in 10 mL of THF and cooled to 0 °C. Butyllithium, 2.5 M in hexanes (3.65 mL, 9.12 mmol), was slowly added, and the solution was stirred in an ice bath for 2 h. Compound **4c** (1.08 g, 0.0035 mol) was dissolved in THF (8 mL) and slowly added to the cold reaction mixture. The mixture was allowed to warm to room temperature for 16 h. A solution of 2.00 g of 36 N H<sub>2</sub>SO<sub>4</sub>, 2.5 mL of H<sub>2</sub>O, and 10.5 mL

of THF was slowly added, and the mixture was refluxed for 6 h before being stirred at room temperature 16 h. The reaction mixture was concentrated and basified with concentrated  $\text{NH}_4\text{OH}-\text{H}_2\text{O}$  (1:1) to pH 10–11 and extracted with  $\text{CH}_2\text{Cl}_2$  (3  $\times$  25 mL). The combined organic extracts were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and concentrated to give 1.23 g of product, which was purified by flash chromatography on silica gel using 10% ( $\text{Et}_2\text{O}-\text{Et}_3\text{N}$ ) (9:1) in 90% hexanes to give 0.39 g of **4c** and 0.71 g (80% adjusted yield) of **4j**. The **4j** was crystallized from petroleum ether to give 0.59 g of fine white crystals: mp 139–141 °C. The product was converted to the HCl salt by adding 1 M HCl in  $\text{Et}_2\text{O}$  to an  $\text{Et}_2\text{O}$  solution of **4j**. Recrystallization from  $\text{MeOH}-\text{Et}_2\text{O}$  gave 0.467 g of **4j**·HCl: mp 278 °C (dec);  $[\alpha]_{\text{D}} -110.2$  °C (*c* 1.00, MeOH).  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  7.66 (m, 2H), 7.43 (m, 3H), 7.21 (d, 1H), 7.12 (s, 1H), 7.00 (d, 1H), 6.20 (s, 1H), 4.30 (m, 1H), 4.16 (m, 1H), 3.98 (m, 1H), 3.75 (m, 1H), 2.93 (s, 3H), 2.04–2.74 (m, 9H). Anal. ( $\text{C}_{24}\text{H}_{27}\text{Cl}_2\text{N}_2\text{O}$ ) C, H, N.

**3 $\alpha$ -(4-Chloro-3-methylphenyl)tropane-2 $\beta$ -carboxylic Acid Methyl Ester (4k) Tosylate.** To  $\text{Ni}(\text{OAc})_2$  (9.21 g, 0.037 mol) in MeOH (50 mL) was added  $\text{NaBH}_4$  (1.4 g, 0.037 mol) in MeOH (20 mL). Compound **8** (2.20 g, 0.0074 mol) in MeOH (20 mL) was added followed by concentrated HCl (3.1 mL, 0.037 mol) in MeOH (5 mL). The black heterogeneous reaction mixture was stirred at reflux for 17 h. The mixture was filtered through Celite, and the filtrate was concentrated in vacuo. The residue was partitioned between  $\text{Et}_2\text{O}$  and concentrated  $\text{NH}_4\text{OH}-\text{H}_2\text{O}$  (1:1). The ether was separated, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated in vacuo to give an orange oil. This oil was chromatographed on silica gel, eluting with hexane– $\text{Et}_2\text{O}-\text{Et}_3\text{N}$  (50:45:5) to afford 0.92 g (46%) of **4k** as a colorless oil. To a solution of **4k** in EtOAc (25 mL) was added 1 equiv of *p*-toluenesulfonic acid in a minimal amount of EtOAc. The resulting solids were separated by filtration and dried to give 1.25 g of **4k**· $\text{CH}_3\text{C}_6\text{H}_4\text{SO}_3\text{H}$  as a white solid: mp 170–171 °C.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , free base)  $\delta$  1.32 (t, *J* = 5.8 Hz, 1H), 1.59 (m, 3H), 2.10 (m, 1H), 2.22 (s, 3H), 2.32 (s, 3H), 2.46 (m, 2H), 3.32 (m, 3H), 3.59 (s, 3H), 6.95 (d, *J* = 3.0 Hz, 1H), 6.98 (s, 1H), 7.25 (d, *J* = 5.5 Hz). Anal. ( $\text{C}_{24}\text{H}_{30}\text{ClNO}_5\cdot 0.25\text{H}_2\text{O}$ ) C, H, N.

**3 $\beta$ -(4-Chlorophenyl)tropane-2 $\beta$ -N-(3'-amino-4,5-dimethylphenyl) Carboxamide (6).** To compound **5** (5.3 g, 0.0189 mol) in  $\text{CH}_2\text{Cl}_2$  (100 mL) was added oxalyl chloride (19.0 mL, 0.0378 mol, 2 M in  $\text{CH}_2\text{Cl}_2$ ). The reaction mixture was stirred at room temperature for 2 h, then concentrated in vacuo. The resulting acid chloride was dissolved in  $\text{CH}_2\text{Cl}_2$  (60 mL) and added to 4,5-dimethyl-1,2-phenylenediamine (6.4 g, 0.0473 mol) in  $\text{CH}_2\text{Cl}_2$  (50 mL). The reaction mixture was stirred, under nitrogen, for a period of 17 h. The solvent was decanted from a gummy residue, and 10%  $\text{NaHCO}_3-\text{CH}_2\text{Cl}_2$  was added. The organic layer was separated, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated in vacuo to give 5.4 g of a foam. This material was chromatographed on silica gel, eluting with EtOAc and then EtOAc–CMA80 (1:1) to afford 3.8 g (51%) of **6** as a yellow amorphous solid.  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.71–1.80 (m, 3H), 2.12 (s, 3H), 2.14 (s, 3H), 2.13–2.39 (m, 3H), 2.39 (s, 3H), 2.66 (d, *J* = 3.0 Hz, 1H), 3.19 (p, *J* = 3.0 Hz, 1H), 3.40–3.60 (bm, 5H), 6.51 (s, 1H), 6.81 (s, 1H), 7.21 (s, 4H).

**3 $\alpha$ -(4-Chloro-3-methylphenyl)-2 $\beta$ -(3'-methyl-1',2',4'-oxadiazol-5-yl)tropane (8).** *n*-Butyllithium (12.0 mL, 0.030 mol, 2.5 M) in hexane was added to 5-bromo-2-chlorotoluene (6.0 g, 0.0292 mol) in THF (50 mL) at –78 °C. The resulting creamy white suspension was stirred for 15 min, and anhydroecgonine oxadiazole (**7**) (3.0 g, 0.0146 mol) in THF (50 mL) was added. The orange reaction mixture was stirred for an additional 3 h, allowing the mixture to come to room temperature. TFA (3.9 mL, 0.050 mol) was added, the mixture was stirred for 15 min and concentrated in vacuo. The resulting residue was treated with concentrated  $\text{NH}_4\text{OH}-\text{H}_2\text{O}$  (1:1) (100 mL) and  $\text{CH}_2\text{Cl}_2$  (100 mL). The organic layer was separated, dried ( $\text{Na}_2\text{SO}_4$ ), and concentrated in vacuo to afford 5.7 g of an orange oil. This oil was chromatographed on silica gel, eluting with ether– $\text{Et}_3\text{N}$

(9:1) to afford 2.0 g (46%) of a mixture that was used without further purification to prepare **4k**.

**Transporter Assays.** The abilities of **2** and its analogues to inhibit uptake of [ $^3\text{H}$ ]dopamine ([ $^3\text{H}$ ]DA), [ $^3\text{H}$ ]serotonin ([ $^3\text{H}$ ]5-HT), or [ $^3\text{H}$ ]norepinephrine ([ $^3\text{H}$ ]NE) by the respective human transporters were evaluated using the appropriate HEK-293 cell line as previously reported.<sup>31</sup>

**Cell Lines and Culture.** Human embryonic kidney (HEK-293) cells stably expressing human DAT, NET, or SERT were maintained as previously described.<sup>31</sup>

Use was made, as previously described,<sup>33</sup> of TE671/RD cells naturally expressing human muscle-type nAChR ( $\alpha 1\beta 1\gamma\delta$ - or  $\alpha 1^*\text{-nAChR}$ ), SH-SY5Y neuroblastoma cells naturally expressing human autonomic  $\alpha 3\beta 4^*\text{-nAChRs}$  (containing  $\alpha 3$ ,  $\beta 4$ , probably  $\alpha 5$ , and sometimes  $\beta 2$  subunits), or transfected SH-EP1 epithelial cells heterologously expressing either human  $\alpha 4\beta 2\text{-nAChR}$ , which is thought to be the most abundant, high affinity nicotine-binding nAChR in mammalian brain, or  $\alpha 4\beta 4\text{-nAChR}$ .<sup>36,37</sup>

**nAChR Functional Assays.** Cells were harvested, seeded onto 24-well plates, and subjected to  $^{86}\text{Rb}^+$  efflux assays as previously described.<sup>33</sup> Specific  $^{86}\text{Rb}^+$  efflux was assessed as the response to a fully efficacious concentration of carbamylcholine alone less than in the presence of efflux buffer alone. Any intrinsic agonist activity of test compounds was normalized, after subtraction of nonspecific efflux, to specific efflux. Antagonism of carbamylcholine-evoked  $^{86}\text{Rb}^+$  efflux was assessed in samples containing the full agonist at a concentration where it stimulates 80–90% of maximal function. For studies of mechanism of antagonism, concentration–response curves were obtained using samples containing the full agonist, carbamylcholine, at the indicated concentrations alone or in the presence of a concentration of the test ligand close to its  $\text{IC}_{50}$  value for inhibition of nAChR function. Ion flux assay results were fit, using Prism (GraphPad), to the Hill equation,  $F = F_{\text{max}}/(1 + (X/Z)^n)$ , where *F* is the test sample specific ion flux as a percentage of control,  $F_{\text{max}}$  is specific ion flux in the absence of test drug (i.e., for control samples), *X* is the test ligand concentration, *Z* is the  $\text{EC}_{50}$  (*n* > 0 for agonists) or  $\text{IC}_{50}$  (*n* < 0 for antagonists), and *n* is the Hill coefficient. All concentration–ion flux response curves were simple and fit well, allowing maximum and minimum ion flux values to be determined by curve fitting, but in cases where antagonists had weak functional potency, minimum ion flux was set at 0% of control. Note that because agonist concentrations used for test ligand antagonism assessments were  $\text{EC}_{80}-\text{EC}_{90}$  values, not all of the data, even at the lowest concentrations of test antagonist, approached 100% of specific efflux as separately determined in sister samples exposed to fully efficacious concentrations of agonist. Note also that it has been repeatedly verified that functional parameters for nicotinic ligands and mechanisms of their action as determined using efflux assays are like those determined using whole-cell current recording techniques.<sup>38</sup>

**Behavior.** All animal experiments were conducted in accordance with the NIH Guide for the Care and Use of Laboratory Animals and Institutional Animal Care and Use Committee guidelines.

**Animals.** Male Institute of Cancer Research (ICR) mice (weighing 20–25 g) obtained from Harlan (Indianapolis, IN) were used throughout the study. Animals were housed in an Association for Assessment and Accreditation of Laboratory Animal Care approved facility, were placed in groups of six, and had free access to food and water. Studies were approved by the Institutional Animal Care and Use Committee of Virginia Commonwealth University.

**Tail-Flick Test.** Antinociception for pain mediated at the spinal level was assessed by the tail-flick method of D'Amour and Smith.<sup>39</sup> In brief, mice were lightly restrained while a radiant heat source was shone onto the upper portion of the tail. To minimize tissue damage, a maximum latency of 10 s was

imposed. Latency to remove the tail from the heat source was recorded for each animal. A control response (2–4 s) was determined for each mouse before treatment, and a test latency was determined after drug administration (nicotine as an analgesic 5 min after subcutaneous administration at 2.5 mg/kg; nicotine administration 15 min after exposure to saline of 3-phenytropane analogue to assess the latter drug's ability to block nicotine-mediated antinociception). Antinociceptive response was calculated as the percentage of maximum possible effect (%MPE), where %MPE = [(test control)/(10 control)] × 100.

**Hot-Plate Test.** Mice were placed in a 10 cm wide glass cylinder on a hot plate (Thermojust Apparatus) maintained at 55 °C for assessment of pain responses mediated at supraspinal levels. To minimize tissue damage, a maximum exposure to the hot plate 40 s was imposed. Measures of control latencies (time until the animal jumped or licked its paws, typically 8–12 s) were done twice for stimuli applied at least 10 min apart for each mouse. Antinociceptive responses after test drug administrations were determined and calculated as the %MPE, where %MPE = [(test latency in s – control latency in s)/(40 s – control latency in s) × 100]. Groups of 8–12 animals were used for each drug condition. Antagonism studies were carried in mice pretreated with either saline or 3-phenytropane analogues 15 min before nicotine. The animals were then tested 5 min after administration of a subcutaneous dose of 2.5 mg/kg nicotine.

**Locomotor Activity.** Mice were placed into individual Omnitech photocell activity cages (28 cm × 16.5 cm; Omnitech Electronics, Columbus, OH) 5 min after subcutaneous administration of either 0.9% saline or nicotine (1.5 mg/kg). Interruptions of the photocell beams (two banks of eight cells each) were then recorded for the next 10 min. Data were expressed as the number of photocell interruptions. Antagonism studies were carried out by pretreating the mice with either saline or 3-phenytropane analogues 15 min before nicotine.

**Body Temperature.** Rectal temperature was measured by a thermistor probe (inserted 24 mm) and digital thermometer (YSI Inc., Yellow Springs, OH). Readings were taken just before and 30 min after subcutaneous injection of either saline or 2.5 mg/kg nicotine. The difference in rectal temperature before and after treatment was calculated for each mouse. The ambient temperature of the laboratory varied from 21 to 24 °C from day to day. Antagonism studies were carried out by pretreating the mice with either saline or 3-phenytropane analogues 15 min before nicotine. The animals were then tested 30 min after administration of a subcutaneous dose of 2.5 mg/kg nicotine.

**Nicotine CPP Assessment.** An unbiased CPP paradigm was utilized in this study as described in Kota et al.<sup>33</sup> Briefly, place-conditioning chambers consisted of two distinct compartments separated by a smaller intermediate compartment with openings that allowed access to either side of the chamber. On day 1, adult male ICR mice were confined to the intermediate compartment for a 5 min habituation period and then allowed to move freely between compartments for 15 min. Time spent in each compartment was recorded. These data were used to separate the animals into groups of approximately equal bias. Days 2–4 were the conditioning days during which the saline group received saline in both compartments and drug groups received sc vehicle or 3-phenytropane analogues 15 min before nicotine (0.5 mg/kg, sc) in one compartment and saline in the opposite compartment for 20 min. Drug-paired compartments were randomized among all groups. Day 5 was the drug free test day, and the procedure was the same as for day 1. Activity counts and time spent on each side were recorded via photosensors using Med Associates interface and software. Separate groups of mice were conditioned with saline or 3-phenytropane analogues alone to investigate if they induce CPP using the same procedure described above. Data were expressed as time spent on drug-paired side minus time spent on saline-paired side. A positive number indicated a preference for the drug-paired side, whereas a negative number indicated an aversion to the drug-paired side. A number at or near zero indicated no preference for either side.

**Acknowledgment.** This work was supported by National Institutes of Health National Cooperative Drug Discovery Group Grant U19 DA019377. Other effort was supported by grants (to R.J.L.) from the National Institutes of Health (Grant DA015389) and the Barrow Neurological Foundation. We thank Lawrence E. Brieady for the synthesis of **4f** and **4k** and Fluvanna Josephson for the synthesis of **4h**, **4i**, and **4j**.

**Supporting Information Available:** Elemental analysis results. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) Centers for Disease Control and Prevention. Smoking and Tobacco Use. Fact Sheet: Health Effects of Cigarette Smoking (updated January 2008). [http://www.cdc.gov/tobacco/data\\_statistics/fact\\_sheets/health\\_effects/effects\\_cig\\_smoking/](http://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/effects_cig_smoking/).
- (2) Centers for Disease Control and Prevention. Annual smoking-attributable mortality, years of potential life lost, and productivity losses—United States. *MMWR Morb. Mortal. Wkly. Rep.* **2005**, *54*, 625–628.
- (3) Benowitz, N. L. Nicotine addiction. *Primary Care: Clin. Office Pract.* **1999**, *26*, 611–631.
- (4) Benowitz, N. L. Clinical pharmacology of nicotine: implications for understanding, preventing, and treating tobacco addiction. *Clin. Pharmacol. Ther.* **2008**, *83*, 531–541.
- (5) Hughes, J. R.; Higgins, S. T.; Bickel, W. K. Nicotine withdrawal versus other drug withdrawal syndromes: similarities and dissimilarities. *Addiction* **1994**, *89*, 1461–1470.
- (6) Tutka, P.; Mosiewicz, J.; Wielosz, M. Pharmacokinetics and metabolism of nicotine. *Pharmacol. Rep.* **2005**, *57*, 143–153.
- (7) West, R.; McNeill, A.; Raw, M. Smoking cessation guidelines for health professionals: an update. Health Education Authority. *Thorax* **2000**, *55*, 987–999.
- (8) Navarro, H. A.; Howard, J. L.; Pollard, G. T.; Carroll, F. I. Positive allosteric modulation of the human cannabinoid (CB) receptor by RTI-371, a selective inhibitor of the dopamine transporter. *Br. J. Pharmacol.* **2009**, *156*, 1178–1184.
- (9) Tutka, P. Nicotinic receptor partial agonists as novel compounds for the treatment of smoking cessation. *Expert Opin. Invest. Drugs* **2008**, *17*, 1473–1485.
- (10) Rollema, H.; Coe, J. W.; Chambers, L. K.; Hurst, R. S.; Stahl, S. M.; Williams, K. E. Rationale, pharmacology and clinical efficacy of partial agonists of alpha4beta2 nACh receptors for smoking cessation. *Trends Pharmacol. Sci.* **2007**, *28*, 316–325.
- (11) Dvoskin, L. P.; Smith, A. M.; Wooters, T. E.; Zhang, Z.; Crooks, P. A.; Bardo, M. T. Nicotinic receptor-based therapeutics and candidates for smoking cessation. *Biochem. Pharmacol.* **2009**, *78*, 732–743.
- (12) Carroll, F. I.; Blough, B.; Abraham, P.; Mills, A. C.; Holleman, J. A.; Wolckenhauer, S. A.; Decker, A. M.; Landavazo, A.; McElroy, K. T.; Navarro, H. A.; Gatch, M. B.; Foster, M. J. Synthesis and biological evaluation of bupropion analogues as potential pharmacotherapies for cocaine addiction. *J. Med. Chem.* **2009**, *52*, 6768–6781.
- (13) Dhillon, S.; Yang, L. P.; Curran, M. P. Bupropion: a review of its use in the management of major depressive disorder. *Drugs* **2008**, *68*, 653–689.
- (14) Ferris, R. M.; Cooper, B. R. Mechanism of antidepressant activity of bupropion. *J. Clin. Psychiatry Monog.* **1993**, *11*, 2–14.
- (15) Nielsen, J. A.; Shannon, N. J.; Bero, L.; Moore, K. E. Effects of acute and chronic bupropion on locomotor activity and dopaminergic neurons. *Pharmacol., Biochem. Behav.* **1986**, *24*, 795–799.
- (16) Nomikos, G. G.; Damsma, G.; Wenkstern, D.; Fibiger, H. C. Effects of chronic bupropion on interstitial concentrations of dopamine in rat nucleus accumbens and striatum. *Neuropsychopharmacology* **1992**, *7*, 7–14.
- (17) Jones, C. N.; Howard, J. L.; McBennett, S. T. Stimulus properties of antidepressants in the rat. *Psychopharmacology (Berlin)* **1980**, *67*, 111–118.
- (18) Lamb, R. J.; Griffiths, R. R. Self-administration in baboons and the discriminative stimulus effects in rats of bupropion, nomifensine, diclofenine and imipramine. *Psychopharmacology (Berlin)* **1990**, *102*, 183–190.
- (19) Ortman, R. The conditioned place preference paradigm in rats: effect of bupropion. *Life Sci.* **1985**, *37*, 2021–2027.
- (20) Tella, S. R.; Ladenheim, B.; Cadet, J. L. Differential regulation of dopamine transporter after chronic self-administration of bupropion and nomifensine. *J. Pharmacol. Exp. Ther.* **1997**, *281*, 508–513.

- (21) Spealman, R. D.; Madras, B. K.; Bergman, J. Effects of cocaine and related drugs in nonhuman primates. II. Stimulant effects on schedule-controlled behavior. *J. Pharmacol. Exp. Ther.* **1989**, *251*, 142–149.
- (22) Runyon, S. P.; Carroll, F. I. Dopamine transporter ligands: recent developments and therapeutic potential. *Curr. Top. Med. Chem.* **2006**, *6*, 1825–1843.
- (23) Runyon, S. P.; Carroll, F. I. Tropane-Based Dopamine Transporter-Uptake Inhibitors. In *Dopamine Transporters: Chemistry, Biology, and Pharmacology*; Trudell, M. L., Izenwasser, S., Eds.; John Wiley & Sons: Hoboken, NJ, 2008; pp 125–169.
- (24) Lerner-Marmarosh, N.; Carroll, F. I.; Abood, L. G. Antagonism of nicotine's action by cocaine analogs. *Life Sci.* **1995**, *56*, PL67–PL70.
- (25) Damaj, M. I.; Slemmer, J. E.; Carroll, F. I.; Martin, B. R. Pharmacological characterization of nicotine's interaction with cocaine and cocaine analogs. *J. Pharmacol. Exp. Ther.* **1999**, *289*, 1229–1236.
- (26) Carroll, F. I.; Pawlusch, N.; Kuhar, M. J.; Pollard, G. T.; Howard, J. L. Synthesis, monoamine transporter binding properties, and behavioral pharmacology of a series of 3β-(substituted phenyl)-2β-(3'-substituted isoxazol-5-yl)tropanes. *J. Med. Chem.* **2004**, *47*, 296–302.
- (27) Kotian, P.; Mascarella, S. W.; Abraham, P.; Lewin, A. H.; Boja, J. W.; Kuhar, M. J.; Carroll, F. I. Synthesis, ligand binding, and quantitative structure–activity relationship study of 3β-(4'-substituted phenyl)-2β-heterocyclic tropanes: evidence for an electrostatic interaction at the 2β-position. *J. Med. Chem.* **1996**, *39*, 2753–2763.
- (28) Carroll, F. I.; Kotian, P.; Dehghani, A.; Gray, J. L.; Kuzemko, M. A.; Parham, K. A.; Abraham, P.; Lewin, A. H.; Boja, J. W.; Kuhar, M. J. Cocaine and 3β-(4'-substituted phenyl)tropane-2β-carboxylic acid ester and amide analogues. New high-affinity and selective compounds for the dopamine transporter. *J. Med. Chem.* **1995**, *38*, 379–388.
- (29) Carroll, F. I.; Mascarella, S. W.; Kuzemko, M. A.; Gao, Y.; Abraham, P.; Lewin, A. H.; Boja, J. W.; Kuhar, M. J. Synthesis, ligand binding, and QSAR (CoMFA and classical) study of 3β-(3'-substituted phenyl)-, 3β-(4'-substituted phenyl)-, and 3β-(3',4'-disubstituted phenyl)tropane-2β-carboxylic acid methyl esters. *J. Med. Chem.* **1994**, *37*, 2865–2873.
- (30) Triggler, D. J.; Kwon, Y. W.; Abraham, P.; Rahman, M. A.; Carroll, F. I. Synthesis of 2-(3-substituted-1,2,4-oxadiazol-5-yl)-8-methyl-8-azabicyclo[3.2.1]octanes and 2 α-(3-substituted-1,2,4-oxadiazol-5-yl)-8-methyl-8-azabicyclo[3.2.1]oct-2-enes as potential muscarinic agonists. *Pharm. Res.* **1992**, *9*, 1474–1479.
- (31) Eshleman, A. J.; Carmolli, M.; Cumbay, M.; Martens, C. R.; Neve, K. A.; Janowsky, A. Characteristics of drug interactions with recombinant biogenic amine transporters expressed in the same cell type. *J. Pharmacol. Exp. Ther.* **1999**, *289*, 877–885.
- (32) Damaj, M. I.; Carroll, F. I.; Eaton, J. B.; Navarro, H. A.; Blough, B. E.; Mirza, S.; Lukas, R. J.; Martin, B. R. Enantioselective effects of hydroxy metabolites of bupropion on behavior and on function of monoamine transporters and nicotinic receptors. *Mol. Pharmacol.* **2004**, *66*, 675–682.
- (33) Carroll, F. I.; Blough, B. E.; Mascarella, S. W.; Navarro, H. A.; Eaton, J. B.; Lukas, R. J.; Damaj, M. I. Synthesis and biological evaluation of bupropion analogues as potential pharmacotherapies for smoking cessation. *J. Med. Chem.* **2010**, *53*, 2204–2214.
- (34) Kota, D.; Martin, B. R.; Robinson, S. E.; Damaj, M. I. Nicotine dependence and reward differ between adolescent and adult male mice. *J. Pharmacol. Exp. Ther.* **2007**, *322*, 399–407.
- (35) Lukas, R. J.; Muresan, A. Z.; Damaj, M. I.; Blough, B. E.; Huang, X.; Navarro, H. A.; Mascarella, S. W.; Eaton, J. B.; Marxer-Miller, S. K.; Carroll, F. I. Synthesis and characterization of in vitro and in vivo profiles of hydroxybupropion analogues: aids to smoking cessation. *J. Med. Chem.* **2010**, *53*, 4731–4748.
- (36) Eaton, J. B.; Peng, J. H.; Schroeder, K. M.; George, A. A.; Fryer, J. D.; Krishnan, C.; Buhlman, L.; Kuo, Y. P.; Steinlein, O.; Lukas, R. J. Characterization of human alpha 4 beta 2-nicotinic acetylcholine receptors stably and heterologously expressed in native nicotinic receptor-null SH-EP1 human epithelial cells. *Mol. Pharmacol.* **2003**, *64*, 1283–1294.
- (37) Gentry, C. L.; Lukas, R. J. Local anesthetics noncompetitively inhibit function of four distinct nicotinic acetylcholine receptor subtypes. *J. Pharmacol. Exp. Ther.* **2001**, *299*, 1038–1048.
- (38) Lukas, R. J.; Fryer, J. D.; Eaton, J. B.; Gentry, C. L. Some Methods for Studies of Nicotinic Acetylcholine Receptor Pharmacology. In *Nicotinic Receptors and the Nervous System*; Levin, E. D., Ed.; CRC Press: Boca Raton, FL, 2002; pp 3–27.
- (39) D'Amour, F. E.; Smith, D. L. A method for determining loss of pain sensation. *J. Pharmacol. Exp. Ther.* **1941**, *72*, 74–79.